

Idaho Statistical Analysis Center

Grants and Research

Idaho State Police

Characteristics and Outcomes of Justice-Involved Youth in Idaho



In collaboration with:

Idaho Department of Juvenile Corrections

Idaho Department of Health and Welfare

Idaho Department of Corrections

Characteristics and Outcomes of Justice-Involved Youth in Idaho

Authors:

Danielle Swerin, M.A.

Thomas Strauss, MPA

For more information:

Email: pgr@isp.idaho.gov

Website: www.isp.idaho.gov/pgr/Research/sac.html

Published: September 2018

This project was supported by Grant No. 2014-BJ-CX-K025 awarded by the Bureau of Justice Statistics. The Bureau of Justice Statistics is a component of the Office of Justice Programs, which also includes the Bureau of Justice Assistance, the National Institute of Justice, the Office of Juvenile Justice and Delinquency Prevention, the Office for Victims of Crime, and the Office of Sex Offender Sentencing, Monitoring, Apprehending, Registering, and Tracking. Points of view or opinions in this document are those of the author and do not necessarily represent the official position or policies of the U.S. Department of Justice, nor the Idaho State Police.

Table of Contents

<i>Executive Summary</i>	1
<i>Review of Research</i>	2
Introduction.....	2
Prevalence of Mental Health and Substance Use Disorders Among Justice-Involved Youth	2
The Impact of Trauma	3
Outcomes	3
Effective Interventions	4
Conclusion	5
<i>Methodology</i>	6
Limitations.....	6
<i>Results</i>	7
Initial Custody Level Assessment (ICLA).....	7
Demographics.....	7
Days in Custody.....	8
Types of Offenses.....	8
Assessments	9
Family Criminality and Abuse.....	9
Massachusetts Youth Screening Instrument (MAYSI).....	10
Youth Level of Service Inventory (YLSI)	11
Progress, Assessment, Reclassification (PAR)	11
Outcomes	12
Employment.....	12
Education	12
Recidivism	13
Youth with 20-511A Court Orders	16
<i>Conclusions and Policy Implications</i>	17
<i>Directions for Future Research</i>	18
<i>References</i>	19
Appendix A: MAYSI Traumatic Experiences Domain	23
Appendix B: Regression Models	23
Appendix C: Female Offenders in IDJC Custody	25
Appendix D: Sex Offenders in IDJC Custody	26

Executive Summary

It is well known that most youth who enter the juvenile justice system struggle with mental health and/or substance abuse concerns. This project sought to discover to what extent this applies to youth under the supervision of the Idaho Department of Juvenile Corrections (IDJC), as well as how these youth perform on a few key outcome measures after leaving custody. To that end, data were collected from multiple state agencies on 843 youth who entered IDJC custody between 2012 and 2016.

Characteristics of Justice-Involved Youth

- 92% suffered from a mental health and/or substance abuse concern.
- Nearly two-thirds had a history of family criminality and/or abuse. Similarly, about two-thirds reported experiencing at least one traumatic event in their lives.
- Youth with mental health concerns alone entered IDJC at an earlier age and stayed in custody longer than any other group.

Outcomes for Justice-Involved Youth

- 56% of youth were charged with a new crime and nearly 1 in 4 were placed under IDOC supervision within three years of release or turning 18.
- The rate of recidivism was substantially higher for youth with substance abuse and co-occurring concerns compared to those without.
- Youth with no mental health or substance abuse concerns were most successful in acquiring jobs after being released from IDJC, while those with a substance abuse concern were the least successful.

20-511A Court Orders*

- 28% of youth in IDJC custody were evaluated by the Department of Behavioral Health (DBH) pursuant to a 20-511A court order. Of these, 99% were formally diagnosed with at least one mental health or substance abuse disorder.
- The top three mental health diagnoses were oppositional-defiant disorder (50%), Attention Deficit Hyperactivity Disorder (ADHD) (47%), and major depression (34%).
- There was not a significant difference in recidivism or employment rates between those with a 20-511A order and those without.

* This section of Idaho Code involves performing mental health assessments and plans of treatment for juvenile offenders when the judge has reason to believe they are suffering a serious emotional disturbance, as defined in section [16-2403](#) of Idaho Code, that has not been adequately addressed.

Review of Research

Introduction

According to 2016 population estimates, there are more than 70 million youth in the United States and more than 400,000 in Idaho alone^{1,2}. One in four Idaho residents are under the age of 18². In Idaho, an estimated 200 per 100,000 juveniles are in residential placement compared to 152 per 100,000 juveniles nationally³. When considering statewide population estimates, this equates to nearly 900 Idaho juveniles. With limited community-based resources, especially for long-term care, the juvenile justice system has become one of the primary mechanisms for accessing mental health and substance abuse services nationally⁴. As a result, a substantial proportion of detained youth in the U.S. and locally suffer from mental health and/or substance use disorders. One meta-analysis found that “adolescents in detention and correctional facilities were about 10 times more likely to suffer from psychosis than the general adolescent population” (p.1010)⁵. In Idaho, more than 58% of detained juveniles meet the criteria for a mental health problem and 42% are positively screened for substance use disorder⁶. Considering the substantial impact of mental health and substance abuse on the juvenile justice system nationally and locally, further inquiry is warranted. Using a solutions-based framework, this section of the report will review prevalence, common underlying factors, short and long-term outcomes, and effective interventions for justice-involved youth with mental health and/or substance abuse concerns.

Prevalence of Mental Health and Substance Use Disorders Among Justice-Involved Youth

It is well-established in research that the majority of justice-involved youth suffer from mental health and/or substance use disorders. One article examining fifteen different studies found that an average of nearly 70% of justice-involved youth have at least one psychiatric disorder⁷. The most common externalizing disorders identified in this study included conduct disorder (46.4%), substance use disorder (45.1%), oppositional defiant disorder (19.8%), and attention-deficit hyperactivity disorder (13.5%). The most common internalizing disorders included anxiety disorder (15.9%), major depression (12%), and post-traumatic stress disorder (9.6%). These results are confirmed by a variety of other studies reporting that an estimated 60-97% of justice-involved youth have at least one mental or behavioral health disorder⁸⁻¹⁴.

In addition to mental health concerns, justice-involved youth also exhibit high rates of substance abuse. According to a nationally representative study of more than 7,000 youth, juveniles in custody are nearly three times more likely to report using marijuana, five times more likely to report using crack or cocaine, and at least four times more likely to report using methamphetamine compared to youth in the general population¹⁵. Research further suggests that approximately 30-65% of justice-involved youth suffer from co-occurring substance abuse and mental health concerns^{8,10,12,16,17}. In fact, the majority (60.8%) of youth with a mental health concern are also diagnosed with substance use disorder¹⁴.

The Impact of Trauma

Considering the pervasiveness of mental health and substance use disorders among justice-involved youth, it is critical to examine potential underlying causes in order to develop and inform effective interventions. Although trauma is frequently included in measurements of mental health (i.e., Posttraumatic Stress Disorder [PTSD]), its unique and substantial impact on risk and outcomes warrants individual examination. Studies of justice-involved youth have consistently found that a history of trauma increases risk for delinquency, mental health problems, substance abuse, suicidal ideation, and PTSD¹⁸⁻²⁶. Some researchers suggest that traumatic experiences may facilitate mental health problems and substance abuse, which subsequently increases risk of delinquency¹⁹. For example, an individual who was abused may develop PTSD and/or begin using substances as a coping mechanism to deal with the trauma symptoms, subsequently increasing risk of delinquency. The high rates of trauma, mental health disorders, and substance abuse among justice-involved youth lends support for this theory. In fact, some studies report almost universal trauma among detained youth, with as many as 95% reporting at least one adverse childhood experience^{15,18,24,27}. From a local perspective, recent analyses suggest that 70% of youth in state custody in Idaho have suffered a history of abuse⁶.

While a history of trauma is prevalent among all justice-involved youth, research suggests that the prevalence and impact of these traumatic experiences may be especially prominent among females. Specifically, multiple studies have identified higher rates of sexual abuse and PTSD among detained females compared to detained males^{10,15,21,24}. Additional research suggests that justice-involved females are more than twice as likely to meet criteria for PTSD compared to males (41% and 18%, respectively)¹⁰ and four to eight times more likely to report a history of sexual abuse^{15,21}. Because of the frequency and impact of trauma on delinquency, especially among females, multiple researchers have highlighted the importance of trauma-informed and gender-specific treatments for justice-involved youth^{4,17,18,21,23,26,28-30}.

Outcomes

In addition to examining potential underlying causes of mental health and substance use disorders, it is also important to assess their impact on both short and long-term outcomes among justice-involved youth. Although mental health and substance use disorders may impact a variety of life outcomes (e.g., education, employment, quality of life), the vast majority of current literature focuses on recidivism. Despite the abundance of available research, results are inconsistent regarding the impact of mental health on reoffending. Some researchers suggest that mental health disorders are associated with an increased risk of recidivism³¹⁻³⁴ while others document no effect^{8,35-37}. Variations based on type of mental health diagnosis have also been noted. For example, a recent meta-analysis found a higher risk of recidivism among juveniles with externalizing disorders (e.g., conduct disorder, ADHD), but not for juveniles with internalizing disorders (e.g., depression, anxiety)³⁴. Some research indicates that internalizing disorders serve as a protective factor against future offending^{32,34}, while other research identifies poorer outcomes among juveniles with anxiety including a higher rate of recidivism among females³⁷.

Contrary to research on mental health alone, it is clear that substance abuse and co-occurring disorders have an impact on delinquency^{8,17,22,31,32,34,35,37,38}. Specifically, substance abuse has been found to increase the risk of recidivism even when accounting for a variety of criminogenic risk factors³⁷. The impact of mental health and substance abuse disorders on recidivism also varies based on gender. Specifically, substance abuse significantly increases the number of offenses for males, but not for females, while having a mental health disorder significantly increases delinquency outcomes for females, but not for males²⁸. In sum, substance use and co-occurring disorders are consistently linked to delinquency outcomes while the impact of mental health alone varies based on gender and type of disorder.

In addition to recidivism, mental health disorders may also impact future employment, education, and financial success²⁹. A study of female detainees found higher levels of employment, educational, and financial problems among participants with a personality disorder compared to participants with no mental health disorder. However, no difference was observed between participants with an Axis I problem (e.g., attention deficit hyperactivity disorder, substance abuse, suicidality, PTSD, dissociation, etc.) and those with no documented mental health concern²⁹.

Effective Interventions

Considering the impact of mental health and substance use disorders on recidivism and other life outcomes, broad and specific evidence-based practices have been identified to improve outcomes among justice-involved youth with behavioral health disorders. Broad, empirically supported strategies consistently identified in available research include:

- (1) prompt and comprehensive screening procedures to identify treatment needs;
- (2) a coordinated system of care that includes cross-agency collaboration; and
- (3) integrated treatment strategies that address mental health, substance use, and criminogenic risk factors^{4,8,10,12,16,25,36,37,39-46}.

The documented benefits of these strategies include improved identification of service needs and likelihood of follow-up treatment, reduction in recidivism, and receipt of community-based mental health care^{40,41,44}. Further research highlights the educational benefits of mental health interventions for justice-involved youth including higher rates of high school completion and lower drop-out rates⁴⁷.

Using this framework, several specific, evidence-based treatment programs have been developed and implemented in communities and detention centers. “The most effective treatment models that have demonstrated delinquency-reducing benefits for youth with mental disorders include Functional Family Therapy [FFT], Treatment Foster Care, and Multisystemic therapy [MST]. Interestingly, all of these therapeutic models are similar in that they involve families and youth, are community based, and deal with problem behaviors and

stresses as a systematic family unit”(p. 7)⁴⁵. In addition to home and community-based models, “Cognitive Behavioral Therapy has emerged as the best validated therapeutic approach for children and adolescents who experience trauma-related symptoms, particularly symptoms associated with anxiety or mood disorders”(p. 43)⁴. Although a variety of other effective models are available, these four interventions* are consistently cited as evidence-based treatment models for justice-involved youth with mental health disorders^{4,17,45,47-49}.

In addition to intensive, community-based treatment models, several interventions have been developed for use within juvenile facilities. With an understanding of the impact of trauma on mental health, substance abuse, and recidivism, effective interventions are frequently developed using a trauma-informed approach. Specifically, researchers have identified four effective trauma-informed treatments for juvenile justice facilities including: Attachment, Self-Regulation, and Competency (ARC); Sanctuary; Structured Psychotherapy for Adolescents Responding to Chronic Stress (SPARCS); and Trauma Affect Regulation: Guide for Education and Therapy (TARGET)²³. Two additional interventions identified by researchers include Trauma and Grief Components Therapy for Adolescents (TGCTA) and Skills Training in Affective and Interpersonal Regulation for Adolescents (STAIR-A)²⁶. Use of these programs in conjunction with the community-based treatments discussed above provide a necessary continuum of care for detained juveniles by implementing well-documented, evidence-based strategies throughout the juvenile justice process*.

Conclusion

It is well-documented in research that a substantial number of justice-involved youth suffer from mental health and/or substance use concerns, which may be manifestations of past trauma. While the individual impact of mental health disorders on recidivism is unclear, it is evident that substance use disorder and a history of trauma increase likelihood of recidivism. Despite the frequency and negative effects of mental health disorders, substance abuse, and past trauma, there are a multitude of evidence-based programs and practices that significantly improve outcomes for justice-involved youth including intensive community-based programs, such as MST and FFT, and trauma-informed interventions within detention centers, such as Sanctuary and TARGET. The implementation of these programs combined with the administration of appropriate screenings/assessments, a wraparound approach to service provision, and integrated treatments that address the complex needs of juvenile offenders are key components of an effective response to justice-involved youth.

* More information about these programs can be found at www.crimesolutions.gov.

Methodology

Characteristics of justice-involved youth in Idaho were measured using data provided by the Idaho Department of Juvenile Corrections (IDJC). Specifically, data included information for all juveniles committed to an IDJC facility between January 1, 2012 through December 31, 2016 (n = 843). Through the use of data sharing partnerships, outcomes were measured using data from a variety of state agencies including the Idaho Department of Labor (IDOL), State Board of Education (SBOE), Department of Behavioral Health (DBH), Idaho Supreme Court (ISC), and Idaho Department of Corrections (IDOC).

Using this data, analyses were conducted for youth with mental health concerns only (MH), substance abuse concerns only (SA), co-occurring concerns (CO), or neither.

- MH = youth in need of mental health treatment while in IDJC custody per the clinician's assessment
- SA = youth in need of substance abuse treatment while in IDJC custody per the clinician's assessment
- CO = youth in need of both mental health and substance abuse treatment while in IDJC custody per the clinician's assessment
- Neither = youth who are *not* in need of mental health or substance abuse treatment per the clinician's assessment.

Limitations

As with any analysis of secondary data, the researchers cannot confirm the data provided were documented consistently or without errors. However, any data entry errors identified during analyses were discussed with the providing agency and modified, if necessary, to ensure accuracy. Due to variations in data collection across facilities and/or changes in policies and procedures over time, some of the variables contained large amounts of missing data. This primarily impacted analyses of Massachusetts Youth Screening Instrument (MAYSI) and Youth Level of Service Inventory (YLSI) scores with 29% of youth missing a MAYSI and 19% missing a YLSI.

Due to changing definitions and methods of data collection, some data were only available for certain years. Specifically, IDOL data was only available for 2016, SBOE data was only available for the 2015 and 2016 school years, and data from the Idaho Supreme Court was only available through 2015. As a result, the sample sizes were much smaller for these analyses. Additionally, the small percentage of juveniles enrolled in post-secondary education prohibited researchers from running predictive analyses with SBOE data.

Because youth with co-occurring concerns accounted for such a large percentage of youth who recidivated as an adult, the other groups had very small sample sizes. As a result, predictive analyses only included the co-occurring group, meaning that researchers were unable to accurately determine if mental health concerns only or substance abuse concerns only had a statistically significant impact on recidivism or employment. While this study did include all youth committed to IDJC within a five-year period, a longer study period could potentially capture a large enough sample to include those other groups.

Results

Initial Custody Level Assessment (ICLA)

Upon arrival at an IDJC facility, youth are assessed by a clinician who determines if the youth needs mental health treatment only (MH), substance use treatment only (SA), both (CO), or neither. Nearly all youth committed to IDJC (92%) were found to have significant enough mental health and/or substance use concerns to warrant treatment. Of those, 42% were identified as having both mental health and substance abuse concerns.

92% of youth in an IDJC facility have a mental health concern, substance abuse concern, or both.


Demographics

The majority of youth committed to IDJC between 2012 and 2016 were white (65%) or Hispanic (21%). Comparatively, 78% of youth with a mental health concern were white and 36% of youth with a substance abuse concern were Hispanic.

Females accounted for 14% of all commitments. Of those, nearly two-thirds (65%) had co-occurring mental health and substance abuse concerns and less than 3% had neither. Comparatively, 37% of males had co-occurring concerns and 9% had neither.

The average age of youth committed to IDJC custody was 16.2 years. Those with a MH concern were among the youngest committed (15.6 years), while those with an SA concern (16.5 years) and co-occurring concerns (16.5 years) were often older. Most youth (86%) were committed during their high school years.

At release, the average youth was 17.9 years old. Individuals with SA (51%) and co-occurring (59%) concerns often remained in custody after turning 18.




Race/Ethnicity, Gender, and Age

Race/Ethnicity	All
White	65%
Hispanic	21%
Black	3%
American Indian	2%
Asian	1%
Pacific Islander	0%
Other/Mixed	1%
Unknown	7%
Gender	
Male	86%
Female	14%
Age at Commitment	
11-13 Years	8%
14-17 Years	86%
18+ Years	7%
<i>Mean</i>	16.2

n = 843

Days in Custody


IDJC has the authority to hold youth on indeterminate sentences, meaning that youth remain in custody until they complete their treatment program or age out of the juvenile justice system. While it is assumed that youth will be released before their nineteenth birthday; they may remain in custody until their twenty-first birthday with approval from the custody review board. The average time spent in custody for all youth who were discharged was 580.80 days (n=747)*. Those with MH concerns tended to have longer terms (664.34 days), while those with SA concerns experienced shorter terms (513.21 days).



Types of Offenses

Property crimes (35%), crimes against persons (34%), and sex crimes (22%) accounted for the majority of all commitments (n=928). However, the most common offense type within ICLA groups varied widely. Sexual offenses were most common in the “neither” (53%) and MH (45%) groups while property crimes were most common among those with SA (55%) and co-occurring (37%) concerns. Drug and alcohol offenses were committed exclusively by those with a documented substance abuse concern (including the co-occurring group).


Most Serious Offense Resulting in Commitment




Assessments

Family Criminality and Abuse

In addition to providing insight on mental health and substance use issues, the ICLA also contains components related to family history (n=822). Most juveniles committed to an IDJC facility between 2012 and 2016 (66%) had a family history of criminal involvement or a documented pattern of abuse, neglect, or probation violations (PVs). These rates are highest among those with SA (68%) and co-occurring (73%) concerns.




In addition to family criminality, the ICLA contains questions regarding where youth were living immediately prior to being committed to an IDJC facility (n=812). The majority of youth (83%) lived in a household that did not include both biological parents. Rates of youth living with one biological parent were highest among those with SA (70%) and co-occurring (66%) concerns.



Massachusetts Youth Screening Instrument (MAYSI)


IDJC uses the Massachusetts Youth Screening Instrument (MAYSI) to further explore the individual needs of youth in custody (n=601). This assessment is conducted soon after a youth enters an IDJC facility and includes an evaluation of suicide risk. Youth can be assigned “caution” status based on the results of this evaluation, and appropriate surveillance measures can be taken to reduce the risk of self-harm while the juvenile is in custody. Higher rates of suicide cautions were issued to youth in the mental health (17%) and co-occurring groups (20%), illustrating the overlap between mental health concerns and suicidal ideation.

Percent of Youth with a Caution or Warning Status for Suicidal Ideation




The Traumatic Experiences domain of the MAYSI gauges an individual’s past proximity to traumatic events, either as a victim or a witness. Nearly two-thirds (65%) of youth reported experiencing at least one traumatic event or trauma symptom, with higher rates occurring in the mental health (66%) and co-occurring (75%) groups (see Appendix A for additional information on how this is measured).

Youth Reporting a History of Trauma




Youth Level of Service Inventory (YLSI)

The Youth Level of Service Inventory (YLSI) is a tool for measuring a youth's criminogenic risks and needs. YLSI assessments are currently conducted at the county level prior to admittance into IDJC custody. The assessment covers seven domains and a total is used to classify individuals as low (0-8 points), moderate (9-22), high (23-34), or very high (35-42) risk. The average total score among all youth was 19.72 (n=682). The mental health group displayed lower total scores with an average of 15.28, while the SA (22.88) and co-occurring (22.31) groups averaged higher scores.



Progress, Assessment, Reclassification (PAR)

IDJC uses the Progress, Assessment, Reclassification (PAR) tool to continually assess risk levels and treatment progress. Youth are re-evaluated every 60 days and assigned a risk level between 1 (lowest) and 5 (highest). At intake, 82% of all youth scored at Level 3 or higher (n=842). On their final evaluations, 81% scored at either Level 1 or 2. The co-occurring group averaged the highest initial and final PAR scores (3.58 and 1.86, respectively), while the neither group averaged the lowest (2.77 and 1.49, respectively).




Outcomes

Employment

Employment data is provided for juveniles who have been out of custody for one year and were at least 17 years old at release. At the time of data collection for this project, IDJC had collected six consecutive quarters of data, beginning with quarter one of 2016. Therefore, employment data were analyzed for juveniles meeting the above criteria who were released between January 1, 2015 and June 30, 2016 (n=166). Overall, 43% of these juveniles had a job with taxable wages within one year of release. Youth with substance abuse concerns had the lowest post-release employment rate (34%) and youth with no mental health or substance abuse concern had the highest (67%).

Post-Release Employment Rates



Top 5 Industries

1. Food Preparation and Serving (31%)
2. Sales (19%)
3. Business/Financial Operations (15%)
4. Construction/Extraction (13%)
5. Farming/Fishing/Forestry (10%)

Education

Of the 239 youth who were released between January 1, 2014 and December 31, 2016 and had their high school diploma or GED at the time of release, 10 (4%) enrolled in college in the 2015 or 2016 school year. Of those who enrolled, most attended a community college in Idaho.

Recidivism

Recidivism was measured using three events as indicators. First, court data was analyzed to determine how many youth had a new criminal case filed against them as an adult. A guilty disposition in any of those cases was used as the second indicator. Court records were available through the end of 2015, yielding a maximum follow-up period of three years for these two indicators (n=201). Finally, data from the Idaho Department of Corrections (IDOC) was used to determine how many youth were under IDOC supervision as an adult*. IDOC data was available through the end of 2017, which expands the maximum follow-up period to five years for this indicator (n=466). At least one year of data was available for all youth included in these analyses.

Within 3 Years:


56% were charged with a new crime

54% received a guilty disposition

24% were under the supervision of the Idaho Department of Corrections

20% were incarcerated in an Idaho Department of Corrections facility

Percent Charged with a New Crime




As illustrated above, youth in the co-occurring group were significantly more likely to be charged with a new crime within three years of release compared to all other groups (see Appendix B for regression results). When considering gender, 100% of females who recidivated were in the co-occurring group (n=11).

*“Under supervision” includes individuals who were on felony probation, committed to an IDOC rider program, committed to a state prison, or on parole.


“Incarcerated” includes individuals who were committed to a state prison or an IDOC rider program.


Month of First Court Filing



Of all youth who were charged with a new crime, the average time between release from IDJC (if 18 years or older at time of release) or turning 18 (if released under 18) and court filing date was 287.73 days. As illustrated above, most reoffending occurred between 4 and 18 months after release or turning 18. A total of 36% were charged with a new crime within one year and 55% within two years. Reoffending rates were highest among youth who were 17 or 18 when first committed to IDJC (62% and 55%, respectively).

New Court Filing by Age at First Commitment






Of the youth who had a criminal court case filed against them as an adult, 70% were charged with a drug or alcohol offense.

Similar to trends found in court records, juveniles with substance abuse or co-occurring concerns were most likely to be placed on IDOC supervision as an adult within five years of being released (if 18 or older at time of release) or turning 18 (if released under 18).

Percent Placed on IDOC Supervision




Youth with 20-511A Court Orders*

During the juvenile court process, the judge may also require youth to be evaluated by the Department of Behavioral Health (DBH) by issuing a 20-511A order. If deemed eligible, these individuals may receive community programming through DBH. Youth who received at least one 20-511A court order comprised 28% of the sample (n=234).

The outcomes of 20-511A evaluations mirrored those of the ICLA assessments. Nearly all youth (96%) were diagnosed with at least one mental illness. Most (86%) were diagnosed with at least two, and 46% were diagnosed with co-occurring mental health and substance abuse problems.

86% of youth who received a 20-511A order had more than one mental health diagnosis and 46% had co-occurring substance abuse disorder.

DBH Diagnoses



Nearly three-quarters (74%) of those evaluated were enrolled in at least one treatment program through DBH. The average number of days each case was open for ongoing treatment was 337.87 days, although nearly half (43%) completed their treatment in less than one year.

Receipt of a 20-511A order as a juvenile did not significantly impact performance on any of the outcome measures (see Appendix B for regression results). Employment rates and court filings were identical between the two groups (43% and 56%, respectively). Additionally, 24% with an order were under IDOC supervision within five years, compared to 27% of those without.

* This section of Idaho Code involves performing mental health assessments and plans of treatment for juvenile offenders when the judge has reason to believe they are suffering a serious emotional disturbance, as defined in section [16-2403](#) of Idaho Code, that has not been adequately addressed.

Conclusions and Policy Implications

Outcomes for justice-involved youth in Idaho are mixed, and often related to the type of problem with which the individual is struggling. On all three recidivism measures, youth with only mental health concerns outperformed youth who struggle with substance abuse. On the other hand, the mental health group performed at the same level or worse than the other groups on the employment and education measures; they held jobs and attended college at equal or lower rates.

In light of these findings, the following recommendations are provided to improve Idaho's juvenile justice system:

- ***Emphasize treatment for youth with substance abuse and co-occurring problems.*** These groups were most likely to recidivate and least likely to sustain employment after release from IDJC. This was particularly true for females considering that 100% of females who were charged with a new crime struggled with co-occurring mental health and substance abuse concerns. The low success rates among these youth may be partially due to their age and length of time spent in custody. Youth with substance abuse and co-occurring concerns spent fewer days in custody and were older when first committed compared to youth with mental health concerns only. Based on these trends, intervening earlier and providing targeted programming for a longer period of time may help improve outcomes for these youth. Because the majority of youth who recidivate are committing drug and alcohol offenses as an adult, it is imperative that programming target the underlying causes of substance abuse specifically.
- ***Provide long-term aftercare services to youth who are released from IDJC.*** While most youth did well during the first three months after release or turning 18, there was a steady increase in new court filings from month 4 to month 18. Although this study did not consider what services are available to youth after release, it is possible that after "aging out" of the juvenile justice system, their access to services is drastically reduced. Partnering with community-based providers to ensure access to services during that 18-month window would enhance the support structure available to struggling youth during this critical time.
- ***Continue to improve collaboration and data sharing among state agencies that have contact with justice-involved youth.*** As noted in the literature review, few studies have attempted to use any measure other than recidivism to determine the effectiveness of the juvenile justice system. Idaho is breaking new ground in collecting information on justice-involved youth from multiple agencies, making it possible to evaluate a variety of outcomes. Using this information, researchers, practitioners, and policymakers can gain a richer understanding of what factors influence their successes and failures as adults. However, at this time, the amount of data available for analysis is limited. Continuing to refine these systems and foster more cooperation between agencies will not only facilitate further research, but as integrated treatment plans become more prevalent, it would aid in the development of this model in Idaho.

Directions for Future Research

This study can be used as a springboard for future research on justice-involved youth in Idaho. The following are examples of research topics that could be explored using this data set or similar data collected at a later date.

- ***Examining the unique characteristics of IDJC's female population.*** Females in IDJC custody are disproportionately affected by co-occurring mental health and substance abuse issues. In fact, every female who recidivated had co-occurring concerns. More in-depth research examining characteristics and outcomes for female offenders could help enhance programming for female offenders specifically.
- ***Further exploration of IDJC's sex offender population.*** The majority of youth with neither mental health nor substance abuse concerns are committed for a sex offense. Sex offenders also represent a relatively large percentage of all juveniles committed to IDJC custody. With this in mind, more in-depth research is needed to examine the unique dynamics of this group.
- ***Identifying factors that influence youth success after release from IDJC custody.*** The present study found that youth perform relatively well for the first three months after release or turning 18. Further examination of what factors are associated with this successful transition could be used to enhance reentry and aftercare programs.
- ***Determining the effect of trauma on characteristics and outcomes of IDJC youth.*** Previous research has indicated that trauma plays an integral role in youth delinquency. However, gaps in data did not allow for the inclusion of trauma in any predictive analyses. Additional data collection would allow this important factor to be included in future research.

References

1. United States Census Bureau. (2016a). *QuickFacts: United States*. Retrieved from <https://www.census.gov/quickfacts/fact/table/US/AGE295216#viewtop>
2. United States Census Bureau. (2016b). *QuickFacts: Idaho*. Retrieved from <https://www.census.gov/quickfacts/fact/table/ID/AGE295216#viewtop>
3. Office of Juvenile Justice and Delinquency Prevention. (2015). *OJJDP statistical briefing book: Juvenile residential placement rates by state, 2013*. Retrieved from <https://www.ojjdp.gov/ojstatbb/corrections/qa08601.asp?qaDate=2013&text=no&maplink=link1>
4. Skowrya, K.R. & Coccozza, J.J. (2007). Blueprint for change: A comprehensive model for the identification and treatment of youth with mental health needs in contact with the juvenile justice system. *National Center for Mental Health and Juvenile Justice*. Retrieved from https://www.ncmhij.com/wp-content/uploads/2013/07/2007_Blueprint-for-Change-Full-Report.pdf
5. Fazel, S., Doll, H., & Langstrom, N. (2008). Mental disorders among adolescents in juvenile detention and correctional facilities: A systematic review and metaregression analysis of 25 surveys. *Journal of the American Academy of Child and Adolescent Psychiatry*, 49(9), 1010-1019.
6. Riley, F. (2014). Idaho's implementation of substance use disorder services for juvenile justice populations. *Corrections Today*. Retrieved from <https://insurancenewsnet.com/oarticle/Idahos-Implementation-of-Substance-Use-Disorder14,-Services-for-Juvenile-Justice-P-a-532498#.WZRIBNKGPRY>.
7. Colins, O., Vermeiren, R., Vreugdenhil, C., van den Brink, W., Doreleijers, T., & Broekaert, E. (2010). Psychiatric disorders in detained male adolescents: A systematic literature review. *Canadian Journal of Psychiatry*, 55(4), 255-263.
8. Schubert, C.A. & Mulvey, E.P. (2014). Behavioral health problems, treatment, and outcomes in serious youthful offenders. *Office of Juvenile Justice and Delinquency Prevention*. Retrieved from <https://www.ojjdp.gov/pubs/242440.pdf>
9. Burke, J.D., Mulvey, E.P., & Schubert, C.A. (2015). Prevalence of mental health problems and service use among first-time juvenile offenders. *Journal of Child and Family Studies*, 24, 3774-3781.
10. Robertson, A.A., Dill, P.L., Husain, J., & Undesser, C. (2004). Prevalence of mental illness and substance abuse disorders among incarcerated juvenile offenders in Mississippi. *Child Psychiatry and Human Development*, 35(1), 55-74.
11. Teplin, L.A., Abram, K.M., McClelland, G.M., Dulcan, M.K., & Mericle, A.A. (2002). Psychiatric disorders in youth in juvenile detention. *Archives of General Psychiatry*, 59(12), 1133-1143.
12. Teplin, L.A., Abram, K.M., McClelland, G.M., Mericle, A.A., Dulcan, M.K., & Washburn, J.J. (2006). Psychiatric disorders of youth in detention. *Office of Juvenile Justice and Delinquency Prevention*. Retrieved from <https://www.ncjrs.gov/pdffiles1/ojjdp/210331.pdf>

13. Drerup, L.C., Croysdale, A., & Hoffman, N.G. (2008). Patterns of behavioral health conditions among adolescents in a juvenile justice system. *Professional Psychology: Research and Practice*, 39(2), 122-128.
14. Schufelt, J.L. & Coccozza, J.J. (2006). Youth with mental health disorders in the juvenile justice system: Results from a multi-state prevalence study. *National Center for Mental Health and Juvenile Justice*. Retrieved from <https://ncmhjj.com/wp-content/uploads/2013/11/PrevalenceRPB.pdf>
15. Sedlak, A.J. & McPherson, K. (2010). *Survey of youth in residential placement: Youth's needs and services*. Retrieved from <https://www.ncjrs.gov/pdffiles1/ojdp/grants/227660.pdf>
16. Hussey, D.L., Drinkard, A.M., & Flannery, D.J. (2007). Comorbid substance use and mental disorders among offending youth. *Journal of Social Work Practice in the Addictions*, 7(1/2), 117-138.
17. Kretschmar, J.M., Butcher, F., Flannery, D.J., & Singer, M.I. (2016). Diverting juvenile justice-involved youth with behavioral health issues from detention: Preliminary findings from Ohio's Behavioral Health Juvenile Justice (BHJJ) initiative. *Criminal Justice Policy Review*, 27(3), 302-325.
18. Becker, S.P. & Kerig, P.K. (2011). Posttraumatic stress symptoms are associated with the frequency and severity of delinquency among detained boys. *Journal of Clinical Child & Adolescent Psychology*, 40(5), 765-771.
19. Bender, K. (2010). Why do some maltreated youth become juvenile offenders? A call for further investigation and adaptation of youth services. *Children and Youth Services Review*, 32, 466-473.
20. Benner, G.J., Stage, S.A., Nelson, J.R., Laederich, M., & Ralston, N.C. (2010). Predicting the cumulative recidivism of juvenile detainees. *Journal of Behavior Analysis of Offender and Victim: Treatment and Prevention*, 2(1), 51-62.
21. Bhatta, M.P., Jefferis, E., Kavadas, A., Alemagno, S.A., & Shaffer-King, P. (2014). Suicidal behaviors among adolescents in juvenile detention: Role of adverse life experiences. *PLOS One*, 9(2), 1-7.
22. Cottle, C.C., Lee, R.J., & Heilbrun, K. (2001). The prediction of criminal recidivism in juveniles: A meta-analysis. *Criminal Justice and Behavior*, 28(3), 367-394.
23. Ford, J.D. & Blaustein, M.E. (2013). Systemic self-regulation: A framework for trauma-informed services in residential juvenile justice programs. *Journal of Family Violence*, 28, 665-677.
24. Ford, J.D., Hartman, J.K., Hawke, J., & Chapman, J.F. (2008). Traumatic victimization, posttraumatic stress disorder, suicidal ideation, and substance abuse risk among juvenile justice-involved youth. *Journal of Child & Adolescent Trauma*, 1, 75-91.
25. Mallett, C.A. (2014). Youthful offending and delinquency: The comorbid impact of maltreatment, mental health problems, and learning disabilities. *Child and Adolescent Social Work Journal*, 31, 369-392.
26. Pickens, I. (2016). Laying the groundwork: Conceptualizing a trauma-informed system of care in juvenile detention. *Journal of Infant, Child, and Adolescent Psychotherapy*, 15(3), 220-230.

27. Abram, K.M., Teplin, L.A., Charles, D.R., Longworth, S.L., McClelland, G.M., & Dulcan, M.K. (2004). Posttraumatic stress disorder and trauma in youth in juvenile detention. *Archives of General Psychiatry*, 61(4), 403-410.
28. Welch-Brewer, C.L., Stoddard-Dare, P., & Mallett, C.A. (2011). Race, substance abuse, and mental health disorders as predictors of juvenile court outcomes: Do they vary by gender? *Child and Adolescent Social Work Journal*, 28, 229-241.
29. van der Molen, E., Vermeiren, R.R.J.M., Krabbendam, A.A., Beekman, A.T.F., Doreleijers, T.A.H., & Jansen, L.M.C. (2013). Detained adolescent females' multiple mental health and adjustment problem outcomes in young adulthood. *Journal of Child Psychology and Psychiatry*, 54(9), 950-957.
30. Greenbaum, C.A. & Javdani, S. (2017). Expressive writing intervention promotes resilience among juvenile justice-involved youth. *Children and Youth Services Review*, 73, 220-229.
31. Feldmann, T.M. (2014). *Factors contributing to juvenile recidivism in a predominately Hispanic population* (Doctoral dissertation). Retrieved from <https://tamucc-ir.tdl.org/tamucc-ir/bitstream/handle/1969.6/558/Taylor%20Feldmann.pdf?sequence=1>.
32. Hein, S., Barbot, B., Square, A., Chapman, J., Geib, C.F., & Grigorenko, E.L. (2017). Violent offending among juveniles: A 7-year longitudinal study of recidivism, desistance, and associations with mental health. *Law and Human Behavior*, 41(3), 273-283.
33. Mallett, C.A. (2013). Factors related to recidivism for youthful offenders. *Social Work Faculty Publications*. Retrieved from http://engagedscholarship.csuohio.edu/cgi/viewcontent.cgi?article=1029&context=clsowo_facpub.
34. Wibbelink, C.J.M., Hoeve, M., Stams, G.J.J.M., & Oort, F.J. (2017). A meta-analysis of the association between mental disorders and juvenile recidivism. *Aggression and Violent Behavior*, 33, 78-90.
35. Elkington, K.S., Teplin, L.A., Abram, K.M., Jakubowski, J.A., Dulcan, M.K., & Welty, L.J. (2015). Psychiatric disorders and violence: A study of delinquent youth after detention. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(4), 302-312.
36. McCormick, S., Peterson-Badali, & Skilling, T.A. (2017). The role of mental health and specific responsivity in juvenile justice rehabilitation. *Law and Human Behavior*, 41(1), 55-67.
37. Schubert, C.A., Mulvey, E.P., & Glasheen, C. (2011). Influence of mental health and substance use problems and criminogenic risk on outcomes in serious juvenile offenders. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50(9), 925-937.
38. Manchak, S.M., Sullivan, C.C., Schweitzer, M., & Sullivan, C.J. (2016). The influence of co-occurring mental health and substance use problems on the effectiveness of juvenile drug courts. *Criminal Justice Policy Review*, 27(3), 247-264.
39. Aalsma, M.C., Brown, J.R., Holloway, E.D., & Ott, M.A. (2014a). Connection to mental health care upon community reentry for detained youth: A qualitative study. *BMC Public Health*, 14(117).

40. Aalsma, M.C., Schwartz, K., & Perkins, A.J. (2014b). A statewide collaboration to initiate mental health screening and assess services for detained youths in Indiana. *American Journal of Public Health, 104*(10), e82-e88.
41. Hammond, S. (2007). Mental health needs of juvenile offenders. *National Conference of State Legislatures*. Retrieved from <https://www.ncsl.org/print/cj/mentaljneeds.pdf>
42. Hodgdon, H. (n.d.). Juvenile offenders and substance use and abuse. *The Future of Children*. Retrieved from https://www.princeton.edu/futureofchildren/publications/highlights/18_02_Highlights_05.pdf
43. Letters, P. & Stathis, S. (2004). A mental health and substance abuse service for a youth detention centre. *Australasian Psychiatry, 12*(2), 126-129.
44. Regional Research Institute for Human Services. (2006). *The integrated co-occurring treatment model (ICT): A new treatment model for youth with co-occurring disorders involved in the juvenile justice system*. Retrieved from <https://www.pathwaysrtc.pdx.edu/pdf/fpS0608.pdf>
45. Underwood, L.A. & Washington, A. (2016). Mental illness and juvenile offenders. *International Journal of Environmental Research and Public Health, 13*, 228-241.
46. Zeola, M.P., Guina, J., & Nahhas, R.W. (2017). Mental health referrals reduce recidivism in first-time juvenile offenders, but how do we determine who is referred? *Psychiatric Quarterly, 88*, 167-183.
47. Cuellar, A. & Dave, D.M. (2016). Causal effects of mental health treatment on education outcomes for youth in the justice system. *Economics of Education Review, 54*, 321-339.
48. Aalsma, M.C., White, L.M., Lau, K.S.L., Perkins, A., Monahan, P., & Grisso, T. (2015). Behavioral health care needs, detention-based care, and criminal recidivism at community reentry from juvenile detention: A multisite survival curve analysis. *American Journal of Public Health, 105*(7), 1372-1378.
49. Zajac, K., Sheidow, A.J., & Davis, M. (2015). Juvenile justice, mental health, and the transition to adulthood: A review of service system involvement and unmet needs in the U.S.. *Children and Youth Services Review, 56*, 139-148.

Appendix A: MAYSI Traumatic Experiences Domain

For the MAYSI, a trained staff member asks juveniles a series of questions to gauge the youth's risk level across a variety of domains. For the traumatic experiences domain, juveniles are asked the following questions:

1. Have you ever in your whole life had something very bad or terrifying happen to you?
2. Have you ever been badly hurt or been in danger of getting badly hurt or killed?
3. Have you ever been raped or been in danger of getting raped?
4. Have you ever had a lot of bad thoughts or dreams about a bad or scary event?
5. Have you ever seen someone severely injured or killed?

Each question is counted as one point towards the final score in the traumatic experiences domain. Therefore, the highest score a youth could receive for this domain is a five. For the purposes of this report, a score of 1-5 was found to indicate that the youth had experienced a traumatic event and/or trauma symptom prior to being committed to an IDJC facility.

Appendix B: Regression Models

Regression models were constructed to determine whether membership in the ICLA Co-Occurring group was a predictor of employment and/or recidivism outcomes. Due to the small nature of the sample size and the composition of the four ICLA groups, predictive analyses were not able to be completed for the MH and SA groups.

Logistic Regression Model: Employment

The employment model for the co-occurring group was not statistically significant ($\chi^2(6) = 7.315$, $p = .293$, $R^2 = .059$). Holding all other variables constant, youth in the co-occurring group were 27% more likely to have a job within one year of release from IDJC than those in the other groups.

Variables	B	Wald	df	p	Odds Ratio
Gender	.829	2.133	1	.144	2.292
Violent Crime as a Juvenile	.425	1.690	1	.194	1.530
Family Criminality	-.422	1.679	1	.195	.656
Living with Biological Parent(s)	-.491	1.208	1	.272	.612
20-511A Order	.095	.062	1	.803	1.099
ICLA Co-Occurring	.239	.472	1	.492	1.270
Constant	-1.052	2.892	1	.089	

Cox Regression Model: New Court Filings

The overall court filing model for the co-occurring group was not statistically significant ($\chi^2 (6) = 11.402, p = .077, -2 \log \text{likelihood} = 1066.63$). Holding all other variables constant, membership in the co-occurring group was associated with a 54% increase in the odds of having a new court filing after becoming at-risk (turning 18 or being released from IDJC, whichever comes last; $p = .032$).

Variables	B	Wald	df	p	Odds Ratio
Gender	0.410	1.584	1	0.208	1.508
20-511A Order	0.086	0.168	1	0.682	1.090
Violent Crime as a Juvenile	-0.321	2.625	1	0.105	0.725
Family Criminality	0.209	1.190	1	0.275	1.233
Living with Biological Parent(s)	-0.275	1.021	1	0.312	0.760
ICLA Co-Occurring	0.430	4.612	1	0.032	1.538

Cox Regression Model: IDOC Supervision

The IDOC supervision model for the co-occurring group was not statistically significant ($\chi^2 (6) = 5.923, p = .432, -2 \log \text{likelihood} = 1357.625$). Holding all other variables constant, membership in the co-occurring group was associated with a 33% increase in the odds of being sentenced to IDOC supervision after becoming at-risk (turning 18 or being released from IDJC, whichever comes last; $p = .133$).

Variables	B	Wald	df	p	Odds Ratio
Gender	0.025	0.008	1	0.930	1.025
20-511A Order	-0.188	0.775	1	0.379	0.829
Violent Crime as a Juvenile	-0.155	0.692	1	0.406	0.856
Family Criminality	0.251	1.870	1	0.171	1.286
Living with Biological Parent(s)	0.081	0.111	1	0.739	1.084
ICLA Co-Occurring	0.287	2.259	1	0.133	1.332




Appendix C: Female Offenders in IDJC Custody

65% of females had co-occurring mental health and substance abuse concerns (compared to 37% of males)

79% of females reported a history of trauma (compared to 62% of males)

48% of females had a new criminal case filed against them within 3 years (compared to 56% of males)

22 was the average YLSI score for female offenders (compared to 19 for males)



Note: Time to recidivism was calculated from the time of release from IDJC custody (if over 18 at release) or from the individual's 18th birthday (if under 18 at release). Court case percentages may not add to 100% due to individuals having one or more cases that involve different types of charges.



Appendix D: Sex Offenders in IDJC Custody

27% of sex offenders had substance abuse concerns
(compared to 77% of non-sex offenders)


693 was the average number of days in custody for sex offenders
(compared to 551 for non-sex offenders)

13 was the average YLSI score for sex offenders
(compared to 22 for non-sex offenders)


33% of sex offenders had a new criminal case filed against them within 3 years
(compared to 58% of non-sex offenders)



Recidivism Measures



Types of Court Cases Filed



Note: Time to recidivism was calculated from the time of release from IDJC custody (if over 18 at release) or from the individual's 18th birthday (if under 18 at release). Court case percentages may not add to 100% due to individuals having one or more cases that involve different types of charges.